

Sex hormones shape acute metabolic response to mild traumatic brain injury: A 3T ¹H-MRS study

Lisa Fuchs^{1,2}, Saskia S. Rusche^{1,3}, Katherine Breedlove², Nico Sollmann^{4,5}, Joachim Strobel⁴, Anja Betz^{1,6,7}, Tesla Shic², Mees Van den Nieuwenhof², Skyler McComas², Alexander P. Lin², Inga K. Koerte^{1,3,6,7,8}

¹ cBRAIN, Department of Child and Adolescent Psychiatry, Psychosomatics and Psychotherapy, University Hospital Ludwig-Maximilians-Universität, Munich, Germany, ² Department of Radiology, Center for Clinical Spectroscopy, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, United States, ³ German Center for Child and Adolescent Health (DZKJ), partner site Munich, Munich, Germany, ⁴ Department of Diagnostic and Interventional Radiology, University Hospital Ulm, Ulm, Germany, ⁵ Department of Diagnostic and Interventional Neuroradiology, School of Medicine, Klinikum rechts der Isar, Technical University of Munich, Munich, Germany, ⁶ Graduate School of Systemic Neurosciences, Ludwig-Maximilians-Universität, Munich, Germany, ⁷ International Max Planck Research School of Translational Psychiatry, Munich, Germany, ⁸ Psychiatry Neuroimaging Laboratory, Department of Psychiatry, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, United States

The author has chosen not to publish their abstract